
Aclopain™

Aceclofenac

Composition:

Aclopain™ Tablet: Each film coated tablet contains Acceclofenac BP 100 mg.

Description:

Aclopain™ is a non-steroidal anti-inflammatory drug (NSAIDs). It works by blocking the action of cyclo-oxygenase, which is involved in the production of prostaglandins. Prostaglandins are produced in response to injury or certain diseases and would cause pain, swelling and inflammation. **Aclopain™** also stimulates cartilage matrix (glycosaminoglycans) synthesis.

Indications:

Aclopain™ is indicated for the relief of pain & inflammation in osteoarthritis, rheumatoid arthritis, ankylosing spondylitis, dental pain, dysmenorrhoea, acute lumbago, knee pain, low back pain.

Dosage & Administration:

Adults:

1 tablet (100 mg) twice daily - 1 tablet in morning & 1 tablet in the evening. Reduce the dosage to 100 mg once daily initially in patient with hepatic dysfunction.

Elderly:

The pharmacokinetics of Acceclofenac is not altered in elderly patients, therefore it is not considered necessary to modify the dose or dose frequency.

Children:

Not recommended.

OR AS DIRECTED BY THE PHYSICIAN.

Side Effects:

Gastrointestinal discomfort, nausea, diarrhea, ulceration, hypersensitivity reactions, rashes, headache, dizziness, nervousness, depression, drowsiness, insomnia, vertigo etc. may occur.

Precautions:

Acceclofenac should be administered with caution to patients with symptoms indicative of gastrointestinal disorders, with a history of peptic ulceration, ulcerative colitis, crohn's disease, hepatic porphyria & coagulation disorders. Patients suffering from severe hepatic & renal impairment must be monitored.

Pregnancy:

This drug should not be used in pregnancy. However, it should be used in pregnancy only if it is clearly needed.

Lactation:

This drug should be avoided during lactation period.

Contraindications:

Acceclofenac is contraindicated in patients with known hypersensitivity to Acceclofenac or aspirin or NSAIDs. It should not be administered to patients with active or suspected peptic ulcer or gastrointestinal bleeding & moderate to severe renal impairment.

Storage Condition:

Store in a cool & dry place, protect from light. Keep out of the reach of children.

Presentation:

Aclopain™ 100 mg Tablet: Each box contains 10x10's tablets in blister pack.

Manufactured by:

Julphar Bangladesh Ltd.

Sreepur, Gazipur, Bangladesh.